

# SAFER AT HOME: PHASE TWO GUIDELINES FOR ALL BUSINESS SECTORS

### PHYSICAL DISTANCING BEST PRACTICES:

- Establish policies and practices for physical distancing between co-workers and between members of the public. (See sector-specific guidelines below for more detailed information on public engagement.)
- Provide clear communication and signage for physical distancing in areas where individuals may congregate, especially at entrances, in seating areas, and in check-out lines.
- Limit the occupancy of physical spaces to ensure that adequate physical distancing may be maintained. (See sector-specific guidelines for more detailed information.)
- Encourage telework whenever possible.
- For those businesses where telework is not feasible, temporarily move or stagger workstations to ensure six feet of separation between co-workers and between members of the public.
- Limit in-person work-related gatherings, including conferences, trade shows, and trainings.
- When in-person meetings need to occur, keep meetings as short as possible, limit the number of employees in attendance, and use physical distancing practices.

### ENHANCED CLEANING AND DISINFECTION BEST PRACTICES:

- ✓ Practice routine cleaning and disinfection of high contact areas and hard surfaces, including check out stations and payment pads, store entrance push/pull pads, door knobs/handles, dining tables/chairs, light switches, handrails, restrooms, floors, and equipment. Follow CDC Reopening Guidance for Cleaning and Disinfection and use an EPA-approved disinfectant to clean. For high contact areas, routinely disinfect surfaces at least every 2 hours. Certain surfaces and objects in public spaces, such as shopping carts and point of sale keypads, should be cleaned and disinfected before each use.
- To the extent tools or equipment must be shared, provide access to and instruct workers to use an <u>EPA-approved disinfectant</u> to clean items before and after use.
- Provide a place for employees and customers to wash hands with soap and water, or provide alcohol-based hand sanitizers containing at least 60% alcohol. (See sectorspecific guidelines for more detailed information.)
- When developing staff schedules, implement additional short breaks to increase the frequency with which staff can wash hands with soap and water. Alternatively, consider providing alcohol-based hand sanitizers with at least 60% alcohol so that workers can frequently sanitize their hands.
- Provide best hygiene practices to employees on a regular basis, including washing hands often with soap and water for at least 20 seconds and practicing respiratory etiquette protocols. A CDC training video is available here: <a href="https://www.cdc.gov/handwashing/videos.html">https://www.cdc.gov/handwashing/videos.html</a>.

### ENHANCED WORKPLACE SAFETY BEST PRACTICES:

✓ Prior to a shift and on days employees are scheduled to work, employers should screen employees prior to starting work. Employees should also self-monitor their symptoms by self-taking of temperature to check for fever and utilizing the questions provided in the <u>VDH Interim Guidance for COVID -19 Daily Screening of Employees</u> before reporting to work. For employers with established occupational health programs, employers can consider measuring temperature and assessing symptoms of employees prior to starting work/before each shift. CDC considers a person to have a fever when he or she has a measured temperature of 100.4° F (38° C) or greater, feels warm to the touch, or gives a history of feeling feverish.

- Implement practices such as those described in <u>VDH Interim Guidance for COVID</u> -19 <u>Daily Screening of Employees</u> for examples of a screening questionnaire. A sample symptom monitoring log is available in this Interim Guidance.
- ✓ Instruct employees who are sick to stay at home and not report to work. If an employee becomes ill or presents signs of illness, follow <u>CDC What to Do if You Are Sick guidance</u>. Employers should post signage in the common languages of the employees telling employees not to come to work when sick.
- Develop or adopt flexible sick leave policies to ensure that sick employees do not report to work. Policies should allow employees to stay home if they are sick with COVID-19, if they need to self-quarantine due to exposure, and if they need to care for a sick family member. Employers should recommend that employees follow <u>CDC</u> guidance on If You Are Sick or Caring For Someone.
- ✓ Some employees are at higher risk for severe illness from COVID-19. These vulnerable employees include individuals over age 65 and those with underlying medical conditions. Vulnerable employees should be encouraged to self-identify and employers should take particular care to reduce their risk of exposure, while making sure to be compliant with relevant Americans with Disabilities Act (ADA) and Age Discrimination in Employment Act (ADEA) regulations.
  - Consider offering vulnerable employees duties that minimize their contact with customers and other employees (e.g., restocking shelves rather than working as a cashier), if agreed to by the employee.
  - Protect employees at <u>higher risk for severe illness</u> by supporting and encouraging options to telework.
  - If implementing health checks, conduct them safely and respectfully, and in accordance with any applicable privacy laws and regulations. Confidentiality should be respected.
  - Other information on civil rights protections for workers related to COVID-19 is available <u>here</u>.
- Designate a staff person to be responsible for responding to COVID-19 concerns.
   Employees should know who this person is and how to contact them.

- Implement staggered shifts for both work periods and break periods. Consider cohort scheduling where groups of employees only work with employees in their group.
- Limit the number of employees in break rooms and stagger breaks to discourage gatherings.
- Use messaging boards or digital messaging for pre-shift meeting information.
- ✓ If the building has not been occupied for the last 7 days, there are additional public health considerations that should be considered, such as taking measures to ensure the <u>safety of your building water system</u>. However, it is not necessary to clean ventilation systems other than routine maintenance as part of reducing the risk of coronavirus transmission.
- Establish a relationship with your local health department and know who to contact for questions.

For healthcare facilities, additional guidance is provided on <u>CDC's Guidelines for</u> Environmental Infection Control in Health-Care Facilities.

### RESOURCES TO PRINT AND DISPLAY:

CDC Symptoms English Spanish CDC Symptoms

CDC Printable Flyer English CDC Printable Flyer Spanish

CDC Printable Flyer Chinese CDC Printable Flyer Korean

CDC Printable Flver Vietnamese FDA information

What Grocery Store and Food Retail Workers Need to Know about COVID-19

CDC Re-Opening America Cleaning and Disinfecting Public Spaces, Workplaces,

Businesses, Schools, and Homes

CDC What You Need to Know About Handwashing VIDEO

## SAFER AT HOME: PHASE TWO GUIDELINES FOR SOCIAL GATHERINGS

#### BEST PRACTICES:

In addition to the best practices in the "Guidelines for All Business Sectors" document, social gatherings should consider the following best practices:

- Post signage at the entrance that states that no one with a fever or symptoms of COVID-19, or known exposure to a COVID-19 case in the prior 14 days, is permitted in the establishment.
- Post signage to provide public health reminders regarding physical distancing, gatherings, options for <u>high-risk individuals</u>, and staying home if sick (samples at bottom of this document).
- Social gatherings should be limited to 50% occupancy of the event space, if applicable, or 50 participants, whichever is less.
- Limit the occupancy of spaces to ensure that six feet of physical distance can be maintained between all organizers, staff, volunteers, and attendees who are not members of the same household.
- Reconfigure seating areas to allow six feet of physical distance between individuals by eliminating or closing select tables or seating areas, or by spreading them out to allow for adequate spacing.
- Ensure exits are configured to reduce the occurrence of bottlenecks at the conclusion
  of the event.
- Practice routine cleaning and disinfection of high contact areas and hard surfaces, including check out stations and payment pads, store entrance push and pull pads, door knobs and handles, dining tables and chairs, light switches, handrails, restrooms, guest lockers, floors, and equipment. Follow CDC Reopening Guidance for Cleaning and Disinfection and use an EPA-approved disinfectant to clean. For high contact areas (e.g., in both public and staff areas), routinely disinfect surfaces at least every 2 hours. Certain surfaces and objects in public spaces, such as point of sale keypads, should be cleaned and disinfected before each use.

- Social gatherings should not include close contact between gathering organizers, staff, and attendees not from the same household. Discontinue activities such as facepainting, temporary tattoo application, or audience participation with performers.
- Events that cannot restrict access to the general public should not take place.
- Where applicable, organizers must create a participant flow plan of modified queue lines to and within the mass gathering. Determine areas likely to become bottlenecks or pinch points and adjust participant flow accordingly.
- ✓ Where applicable, install visible markers for queue lines that separate people by six feet of physical distance. Provide physical guides to seating areas including floor decals, colored tape, or signs to indicate where attendees should not sit, stand, or congregate. Sample markers are available in the VDH Business Toolkit.
- Do not include side events as part of the gathering. These are events associated with but located outside of the formal gathering location, such as souvenir vending stations, gaming areas, petting zoos, and designated food truck areas.
- Designate a coordinator who will be responsible for identifying COVID-19 related issues and their impact on the gathering, including coordinating health preparedness for the event and emergency response planning.
- Develop an emergency and medical response plan for COVID-19. This plan should include information about how attendees should interface with the local healthcare system (who to speak with or call if they feel sick) and procedures for isolating sick attendees and reporting illnesses to event organizers. The response plan must also include a plan for organizers to notify the local health department and all participants if any exposure to COVID-19 occurs at the event.
- Develop a mitigation plan for COVID-19, to include how all requirements listed here will be met, and how guidelines and requirements will be enforced at the gathering.
- Consider limiting the duration of the gathering, especially for indoor gatherings and gatherings involving activities that would increase respiration (e.g., cheering, singing).
- Consider delaying gatherings where the target demographic is persons in high risk populations.

- Use technology solutions where possible to reduce person-to-person interaction, including contactless payment and ticket scanning.
- ✓ Install physical barriers, such as sneeze guards and partitions, at staffed kiosks.
- All bathrooms, hand-washing stations, and portable toilets should be placed in a convenient location for use by attendees but away from congregate areas and arranged in a configuration that maintains at least six feet of physical distance.
- Provide a place for staff and attendees to wash hands with soap and water, or provide alcohol-based hand sanitizers containing at least 60% alcohol.
- Where applicable, provide best hygiene practices to employees on a regular basis, including washing hands often with soap and water for at least 20 seconds and practicing respiratory etiquette protocols. A CDC training video is available here: <a href="https://www.cdc.gov/handwashing/videos.html">https://www.cdc.gov/handwashing/videos.html</a>.
- Close off areas used by a sick person and do not use them until after cleaning and disinfection. Wait 24 hours before cleaning and disinfecting. If it is not possible to wait 24 hours, wait as long as possible. Ensure safe and correct application of disinfectants and keep disinfectant products away from children.
- Attendees should leave as quickly as possible following the event in order to limit contact and exposure to themselves and other attendees.
- Establish relationships with key community partners and stakeholders, such as the local health department, community leaders, hospitals, and law enforcement.
   Collaborate and coordinate with them on broader planning efforts for the gathering.
- ✓ Identify actions to take if the event needs to be postponed or cancelled.
- Private events may be held at venues provided they can comply with these guidelines.

### RESOURCES TO PRINT AND DISPLAY:

CDC Symptoms English CDC Symptoms Spanish

CDC Printable Flyer English CDC Printable Flyer Spanish

### CDC Printable Flyer Chinese

### CDC Printable Flyer Korean

CDC Printable Flyer Vietnamese

FDA information